

ENTERIC OUTBREAK MANAGEMENT CHECKLIST

For Congregate Living Settings

Refer to this checklist to manage outbreaks as per the Ministry of Health's protocols and the Windsor-Essex County Health Unit (WECHU) recommendations. Retain for your records.

Setting Name:		Outbreak #: 2268 – –		Date:		
Outbreak Declaration: Suspect Confirmed						
Affected Area: Entire living setting □ OR Name of unit(s)/floors(s):						
Case definition: determined by the WECHU (Click here or visit wechu.org)						
☐ Abnormal temperature ☐ Vo		omiting	□ Dia	arrhea		
☐ Chills ☐ Cra		amps	□ Na	usea		
☐ Malaise/fatigue ☐ He		eadache	□ Ot	her:		
CONTACT						
Identify the designated WECHU nurse for your outbreak: Nurse Name: Phone #: 519-258-2146 ext For any questions or concerns please contact your designated nurse or the Infectious Disease Prevention (IDP) department at 519-258-2146 Ext 1420. The WECHU business hours are from 8:30am - 4:30pm Monday to Friday. Contact the After-Hours hotline at 519-973-4510 to speak with on-call personnel outside of WECHU business hours.						
IMMEDIATE ACTIONS						
If someone residing	Individual should remain in their room.					
in the setting is symptomatic:	Implement additional precautions (i.e., contact/droplet), if possible.					
*More information	Provided the necessary medical assessments.					
available below Test to determine specific illness.						
		TESTING				
Ensure there is a plan in place to test individuals residing in the setting for pathogens in accordance with internal policies and procedures. Refer to the PHO's <u>Kit and Test Ordering Instructions</u> webpage for more information on enteric illness testing.						
Consult with your WECHU outbreak nurse and a primary care provider for more information on specific test types and lab requisitions.						
LINE LISTS						
□ wechu.org).	Create a line list of ill individuals who belong to the outbreak (<u>click here</u> to download the line list or visit wechu.org). *Only include those on the line list who meet case definition (see above).					
☐ Update and fax line lists <u>daily</u> to WECHU <u>by 10:00 am</u> to fax (519)-977-5097.						
COMMUNICATION						
Post outbreak signage at all entrances of building.						
Notify all staff, students, volunteers, families, and visitors of the outbreak. The WECHU will send your setting an Advisory Notice to reflect the current outbreak.						
An Outbreak Notification will be <u>posted on the WECHU website</u> alerting others of a current outbreak congregate living setting, however identifying details such as the name of the setting and the address not be disclosed.						



ENTERIC OUTBREAK MANAGEMENT CHECKLIST

For Congregate Living Settings

	Convene an Outbreak Management Team (OMT) and meet daily to review the status of the outbreak and communicate updates to the WECHU.						
PUBLIC HEALTH INSPECTOR							
	PHI Name:	designated Public Health Inspector (PHI) from WECHU for your setting, if applicable: Phone #: 519-258-2146 ext lealth Inspector (PHI) may reach out to conduct a site visit.					
	IPAC OUTBREAK MEASURES						
	Refer to WECHU IPAC Hub website and the Ministry of Health documents for additional resources related to outbreak control measures: Recommendations for Outbreak Prevention and Control in Institutions and Congregate Living Settings – February 2025, or as current. Appendix 1: Ontario Public Health Standards, Gastroenteritis Outbreaks in Institutions and Public Hospitals – May 2022, or as current.						
	Outbreak Education	Provide education to those in the setting about outbreak control measures, such as proper use of masks, cleaning hands, and reporting new symptoms to staff.					
	Case Control Measures	Symptomatic individuals should remain in their room and cohort cases (i.e., limit movement of individuals between outbreak areas and non-outbreak areas) where/when possible. Refer to Section 7.3 of the Infection Prevention and Control (IPAC) and Outbreak					
		Management in Congregate Living Settings document for more information on outbreak control measures by outbreak type.					
	Additional Precautions	All positive cases should be placed on contact precautions in addition to routine practices. Droplet precautions may be required based on PHO's <u>Risk Assessment</u> <u>Related to Routine Practices and Additional Precautions</u> . Refer to <u>Section 2.0</u> of the <u>Infection Prevention and Control (IPAC) and Outbreak Management in Congregate Living Settings document or <u>PHO's Routine and Additional Precautions</u> webpage for more information.</u>					
		Post additional precautions signage on the door of case rooms, when possible.					
	Staff/ Students/ Volunteers Control Measures	Minimize movement of staff/students/volunteers between outbreak and non-outbreak areas as much as possible (i.e., cohorting staff).					
		Exclude ill staff/students/volunteers for at least 48 hours after their last symptom or longer if indicated by internal policies. NOTE: If a specific illness is known, disease-specific exclusions apply.					
		If dietary staff become ill while working, discard all ready-to-eat food they prepared while on shift.					
	Dietary	Ensure there is a dining policy in place to separate exposed and unexposed individuals.					
		Provide in-room tray service meals within the outbreak area, if available. Ensure the staff who deliver meals are practicing proper hand hygiene in between rooms.					
		DO NOT dispose of food samples until speaking with your PHI or WECHU nurse.					
	Visitor Control Measures	Restrict visitors to essential caregivers on affected units.					
		Ensure those who do visit:Are screened for signs and symptoms of illness.					



ENTERIC OUTBREAK MANAGEMENT CHECKLIST

For Congregate Living Settings

		Practice vigilant hand hygiene.				
		Visit individuals in their rooms and				
		Visit only one individual; do not mingle with others.				
		Use appropriate PPE especially if providing direct care.				
		Ill visitors should be advised not to visit white ended.	le they are ill and wait until symptoms have			
	Increased	For more information, refer <u>Section 5.1</u> of the <i>Infection Prevention and Control (IPAC)</i>				
	Environmental	and Outbreak Management in Congregate Living Settings document or to PHO's Best				
	Cleaning	Practices for Environmental Cleaning – Apri				
	Hand Hygiene	Ensure proper handwashing is maintained by everyone in the setting by providing ample supply of soap and 70-90% alcohol-based hand sanitizers (e.g., where sinks are not readily available).				
		Ensure proper PPE, for example, masks (N95 where applicable), gloves, gowns and eye protection, are available and accessible throughout the setting.				
	PPE	Provide a container for soiled PPE/linen:				
		 If the container is located <i>inside</i> the individual's room, the container must be a minimum of 6ft or more away from the individual's bed. If not possible, place the container <i>outside</i> the room a minimum of 6ft away from any clean linen. 				
		*Ensure alcohol-based hand sanitizer is available by the container.				
		Refer to <u>Section 2.0</u> and <u>Section 7.3</u> of the <i>Infection Prevention and Control (IPAC) and Outbreak Management in Congregate Living Settings</i> document for more information on the required PPE for a respiratory outbreak.				
	Audit	Increase audits of staff practices (e.g. hand hygiene, cleaning, use of PPE, etc.).				
		Reschedule communal activities in the affected area.				
	Activities	Exceptions regarding non-outbreak units/floors should be discussed with the OMT involving outside groups such as entertainers, volunteer organizations, and community groups.				
	Admissions/ Readmissions & Transfers	Limit, if possible, when a new outbreak has been declared. For specific guidance on admissions/readmission, refer to Section 8.5 and 8.6 (page 73-74) of the Recommendations for Outbreak Prevention and Control in Institutions and Congregate Living Settings – February 2025 or as current.				
	Medical/Other Appointments	If possible, reschedule non-urgent appointments until the outbreak is over.				
_	nature and ignation:		Date:			