519-258-2146 | wechu.org |







1005 Ouellette Avenue, Windsor, ON N9A 4J8 | 33 Princess Street, Leamington, ON N8H 5C5

HIGH RISK MENINGOCOCCAL VACCINE ORDER FORM

FAX: (519) 977-1711 | EMAIL: vaccine@wechu.org

Date:	Phone Number:
Facility Name:	
Contact Person:	Fax Number:
Pick Up by: Medex □ Staff □ S	Site: Windsor □ Leamington □
Eligibility:	
1. Functional or anatomic asplenia	
 Complement, properdin, factor D of Cochlear implants (pre/post implants) Acquired complement deficiencies HIV 	nt) (e.g. receiving eculizumab)
 Complement, properdin, factor D of 3. Cochlear implants (pre/post implant 4. Acquired complement deficiencies 5. HIV Vaccines 	nt) (e.g. receiving eculizumab) # of doses
Complement, properdin, factor D of 3. Cochlear implants (pre/post implant 4. Acquired complement deficiencies 5. HIV Vaccines Men-C-ACYW (Menactra/Nimenrix/Menve	# of doses (e.g. receiving eculizumab) # of doses
 Complement, properdin, factor D of 3. Cochlear implants (pre/post implant 4. Acquired complement deficiencies 5. HIV Vaccines	# of doses (e.g. receiving eculizumab) # of doses
Complement, properdin, factor D of 3. Cochlear implants (pre/post implant 4. Acquired complement deficiencies 5. HIV Vaccines Men-C-ACYW (Menactra/Nimenrix/Menve	# of doses (e.g. receiving eculizumab) # of doses
 Complement, properdin, factor D of 3. Cochlear implants (pre/post implant 4. Acquired complement deficiencies 5. HIV Vaccines Men-C-ACYW (Menactra/Nimenrix/Menve 4C MenB (Bexsero) - (age 2 months – 17 y 	# of doses (e.g. receiving eculizumab) # of doses
2. Complement, properdin, factor D of 3. Cochlear implants (pre/post implant 4. Acquired complement deficiencies 5. HIV Vaccines Men-C-ACYW (Menactra/Nimenrix/Menve 4C MenB (Bexsero) - (age 2 months – 17 y Client Information	# of doses (e.g. receiving eculizumab) # of doses

For more details and recommended number of doses by age, please refer to the Publicly Funded Immunization Schedule for Ontario

Vaccine orders will only be processed when accompanied by the most current 4 week temperature log.

Coolers must be pre-chilled between 2-8° C

All transport equipment must be present to receive vaccine.